

Size-Exclusion Chromatography-based isolation minimally alters Extracellular Vesicles' characteristics compared to precipitating agents

Ana Gámez-Valero^{1,2}, Marta Monguió-Tortajada¹, Laura Carreras-Planella¹, Marcel·la Franquesa¹, Katrin Beyer², Francesc E. Borràs^{1,3*}

¹REMAR-IVECAT Group, Health Science Research Institute Germans Trias i Pujol, Can Ruti Campus, Badalona, Spain;


²Department of Pathology, Hospital Universitari and Health Sciences Research Institute Germans Trias i Pujol, Universitat Autònoma de Barcelona, Spain;

³Nephrology Service, Germans Trias i Pujol University Hospital, Badalona, Spain

Key words: PROSPR, Size Exclusion Chromatography, PEG precipitation, EVs isolation, exosomes

*Correspondence should be addressed to:

F.E.B.: feborras@igtp.cat, Tel: 0034 93 497 86 71; Fax: 0034 93 497 86 68



Supplementary Figure 1. Representative flow cytometry plots of cells incubated for 24h at 37°C with EVs isolated from stimulated THP-1 cells by SEC, PEG or PROSPR methods. Cells exposed to PBS were used as control condition. Viable cells were gated according to their FSC-A/SSC-A profiles and dead cells were gated by 7AAD⁺ staining. One representative experiment for THP-1 monocytes and for macrophages are shown.